

RESTORATION STRATEGY / FRAMEWORK FOR SUSTAINABLE RECOVERY AND CONSERVATION OF THE NABAKAZI WETLAND ECOSYSTEM



March, 2026

A publication of:

**National Association of Professional Environmentalists
(NAPE)**

With support from:

Partner für Klima und Natur gGmbH (PKN).



**PARTNER FÜR
KLIMA UND NATUR**

Table of Contents

Table of Contents.....	2
1.0 Executive Summary.....	3
2.0 Acknowledgements.....	4
3.0 Preamble.....	5
4.0 Overview of Nabakazi Wetland Ecosystem.....	6
4.1 Geographic and Hydrological Context.....	6
4.2 Ecological Characterization.....	7
4.3 Socio-Economic Significance.....	9
5.0 Justification for the Restoration Framework.....	11
5.1 Degradation Status and Drivers.....	11
5.2 Policy and Legal Justification.....	12
6.0 Restoration Framework (Five Stages).....	13
6.1 Stage 1: Area Assessment	13
6.2 Stage 2: Planning and Design.....	14
6.3 Stage 3: Implementation of Interventions.....	14
6.4 Stage 4: Monitoring of Results.....	15
6.5 Stage 5: Post-Restoration Management (Adaptive Management).....	15
7.0 Institutional Arrangements and Stakeholder Roles.....	16
7.1 District and Local Government Authorities.....	16
7.2 National Association of Professional Environmentalists (NAPE) Technical Team....	16
7.3 Community Wetland Management Committees (CWMCs).....	17
7.4 Government Agencies and Sectoral Partners.....	17
7.5 Private Sector and Civil Society Partners.....	17
7.6 Coordination	17
8.0 Implementation Strategy and Operational Plan.....	18
8.1 Phased Implementation Approach.....	18
8.2 Community Engagement and Participation.....	19
8.3 Resource Mobilization and Management	19
8.4 Timelines and Work Plan.....	19
8.5 Monitoring, Evaluation, and Learning.....	19
8.6 IMPLEMENTATION SCHEDULE AND RESOURCE PLAN.....	19
8.7 Risk Management.....	22
9.0 Monitoring, Evaluation, and Adaptive Management.....	23
9.1 Objectives of Monitoring and Evaluation.....	23
9.2 Monitoring Indicators.....	24
9.3 Monitoring Methods.....	25
9.4 Evaluation and Reporting.....	25
9.5 Adaptive Management.....	25
10. Sustainability Measures and Long-Term Management.....	26
10.1 Ecological Sustainability.....	26
10.2 Community and Livelihood Sustainability	27
10.3 Institutional and Governance Sustainability.....	27
10.4 Financial Sustainability.....	28
10.5 Knowledge Management and Learning.....	28
11.0 References	29

1.0 Executive Summary

The Nabakazi Wetland Ecosystem spans Mubende, Kassanda and Gomba, districts in Central Uganda and functions as a critical ecological and hydrological resource. It is a headwater for the Katonga, Kafu, Nkuse, and Muzizi river systems, which feed into Lake Victoria, Lake Kyoga, Lake George, and Lake Albert, and forms a significant component of the greater Nile Basin. The wetland provides a multitude of ecosystem services including water purification, groundwater recharge, flood attenuation, carbon sequestration, and biodiversity conservation.

Recent assessments have highlighted the extensive degradation of the wetland due to agricultural encroachment, deforestation, soil erosion, invasive species, and unregulated human activities. These pressures have reduced the wetland's ecological integrity and compromised its ability to provide essential services to the local communities.

The Nabakazi Wetland Restoration Framework provides a participatory, evidence-based, and technically grounded approach to reversing these trends. The framework is structured around five stages: first, a comprehensive area assessment establishes ecological, hydro-

logical, and socio-economic baselines. Second, planning and design translates these assessments into a detailed restoration plan, incorporating stakeholder perspectives. Third, implementation of interventions includes ecological rehabilitation, hydrological restoration, invasive species management, and livelihood enhancement. Fourth, rigorous monitoring of results ensures accountability and guides adaptive management. Fifth, post-restoration management embeds long-term governance, institutionalization, and sustainability strategies.

This framework is fully aligned with national policies, including the Uganda Wetlands Policy (1995) and the National Environment Act (2019), as well as international instruments such as the Ramsar Convention, the African-Eurasian Waterbird Agreement, and the Convention on Biological Diversity. By integrating ecological restoration with socio-economic development and governance strengthening, the framework promotes both environmental sustainability and community resilience.

2.0 Acknowledgements

The preparation of this framework benefited immensely from the collaboration of multiple stakeholders. The Mubende District Environment and Natural Resources Office led the coordination, field mapping and peer reviews while the National Association of Professional Environmentalists (NAPE) led the field mobilization, provided technical guidance, executed baseline studies, and developed the restoration methodology. Financial and technical support was provided by Partner Fur Klima und Natur (PKN), which enabled the execution of field activities, design, production and printing of the final product.

Local communities and leaders from Madudu, Butologo, Kitenga, Kalonga, and Bwaka-go played a central role in participatory assessments, validation workshops, and practical planning exercises. Additional support was provided by NEMA, MWE, civil society organizations, academic institutions, private sector developers among others. The collective contributions of these stakeholders ensured that the framework is practical, scientifically sound, and responsive to local ecological and socio-economic realities. Their participation has laid the foundation for sustainable restoration of the Nabakazi Wetland Ecosystem.

3.0 Preamble

The Nabakazi Wetland Restoration Framework was developed as a direct response to the escalating ecological degradation threatening one of Central Uganda's most vital wetland ecosystems. Over recent years, the wetland has faced significant pressure from human activities, including unsustainable agricultural expansion, deforestation of riparian buffers, encroachment into critical wetland zones, and the proliferation of invasive plant species. These disturbances have undermined the wetland's ecological integrity, reduced its capacity to provide essential ecosystem services, and increased vulnerability of surrounding communities to climate variability, flooding, and water scarcity.

The overarching objective of the framework is to provide a structured, participatory, and scientifically informed approach to restore the ecological functions of the Nabakazi Wetland, while simultaneously improving local livelihoods, enhancing household food security, and building resilience to climate-related risks. By aligning ecological restoration efforts with the socio-economic needs of the wetland-dependent communities, the framework ensures that conservation measures are practical, equitable, and sustainable.

Development of the framework involved a comprehensive, multi-level consultative process. Field-based assessments were conducted across key subcounties, complemented by stakeholder workshops, focus group discussions with local resource users, and technical peer reviews by experts in wetland ecology and restoration. These participatory processes ensured that the framework reflects local realities, leverages indigenous knowledge, and incorporates the priorities of both communities and government stakeholders.

The methodology underpinning the framework is grounded in internationally recognized standards for ecological restoration, drawing from the Society for Ecological Restoration (SER), the International Union for Conservation of Nature (IUCN), and the Food and Agriculture Organization (FAO). It emphasizes adaptive management, allowing restoration strategies to be continuously refined based on monitoring results, ecological feedback, and evolving socio-economic contexts. Stakeholder engagement is central to the approach, ensuring active community participation in planning, implementation, and long-term management of restored areas. Furthermore, the framework integrates ecological restoration with socio-economic development interventions, thereby promoting a holistic approach that is both environmentally effective and socially inclusive. By combining rigorous scientific principles with participatory governance and community-centered livelihood strategies, the framework provides a scalable, replicable, and sustainable model for wetland restoration in Central Uganda and similar ecological landscapes.

4.0 Overview of Nabakazi Wetland Ecosystem

4.1 Geographic and Hydrological Context

The Nabakazi Wetland Ecosystem is one of Central Uganda's most extensive and ecologically significant wetlands. While the majority of its area lies within Mubende District, the wetland extends into Kassanda and Gomba, districts, forming a trans-district ecological landscape that sustains both biodiversity and human livelihoods. Covering approximately 64 square kilometers, the wetland originates in Madudu Subcounty, an area characterized by undulating highlands and seasonal streams that serve as the primary sources of water for the wetland. From Madudu, the wetland flows through a series of interconnected sub counties—including, Kitenga, Kalonga, and Bwakago—traversing diverse landscapes, ranging from gently sloping farmlands to riparian forests and marshy floodplains. This physical connectivity ensures the continuity of ecological processes and sustains a mosaic of habitats critical for wetland-dependent species.

Hydrologically, the Nabakazi Wetland is a vital node in Uganda's inland water system, contributing water to four major catchments: Katonga, Kafu, Nkuse, and Muzizi. Through these networks, the wetland's waters eventually reach major water bodies such as Lake Victoria, Lake Kyoga, Lake George, and Lake Albert, and ultimately form part of the Nile River Basin. This connection highlights the wetland's regional importance as a source of freshwater, sediment regulation, and nutrient transport. The wetland's natural hydrological functions include flood attenuation, water storage, sediment trapping, and groundwater recharge. During periods of heavy rainfall, the wetland absorbs excess water, reducing downstream flood risks and protecting settlements, farmlands, and infrastructure. In dry

periods, it sustains base flow in rivers, maintains wetland vegetation, and replenishes groundwater reserves that local communities rely on for domestic use and irrigation.

The Nabakazi Wetland also plays a critical role in controlling soil erosion and sedimentation. Its dense vegetation, particularly papyrus (***Cyperus papyrus***), reeds (***Phragmites mauritianus***), and sedges, stabilizes riverbanks and filters suspended sediments and pollutants before they enter downstream water bodies. These processes help maintain water quality in the rivers and lakes it feeds, benefiting both aquatic life and human populations. Additionally, by acting as a natural water reservoir, the wetland contributes to local microclimatic regulation, reducing temperature extremes, mitigating the impacts of droughts, and sustaining agricultural productivity in the surrounding sub counties.

Socio-ecologically, the wetland is indispensable to the communities that live along its margins. Agricultural lands adjacent to the wetland depend on the nutrient-rich soils deposited by seasonal flooding. Farmers grow coffee, maize, bananas, and other crops that form the backbone of local livelihoods. The wetland also supports artisanal fishing, harvesting of papyrus for craft-making, and provision of grazing grounds for livestock during dry periods. Its ecological health is therefore directly linked to community food security, income generation, and resilience to climate variability.


However, the strategic location and ecological importance of the wetland also


make it highly vulnerable. Encroachment for agriculture, uncontrolled clearing of riparian vegetation, settlement expansion, and unsustainable extraction of resources threaten its hydrological functions and ecosystem services. Loss of vegetation along the wetland margins reduces its ability to filter sediments, regulate water flow, and provide habitats for fish, amphibians, and birds—including the Grey Crowned Crane (***Balearica regulorum***), Uganda's national bird. The cumulative effects of these pressures compromise not only local livelihoods but also downstream water security, aquatic biodiversity, and the

integrity of Uganda's interconnected lake and river systems.

In conclusion, the Nabakazi Wetland is more than just a local natural resource; it is a regional ecological linchpin that sustains hydrological, ecological, and socio-economic functions across multiple districts. Its protection and restoration are essential for maintaining water availability, soil fertility, biodiversity conservation, and the livelihoods of tens of thousands of residents, as well as for safeguarding the health of the broader Nile Basin.

MAP SHOWING NABAKAZI CATCHMENT AREA





4.2 Ecological Characterization

The Nabakazi Wetland Ecosystem is characterized by a rich mosaic of habitats, which together support a high diversity of plant and animal species, providing critical ecological functions that sustain both local livelihoods and broader environmental processes. Its ecological composition is influenced by hydrology, soil types, elevation gradients, and human activity, creating a dynamic system of marshes, riparian forests, open water channels, and grasslands.

The wetland's dominant vegetation includes extensive stands of papyrus (***Cyperus papyrus***), reeds (***Phragmites mauritianus***), sedges, and other hydrophilic plants that thrive in saturated soils. Papyrus swamps form dense mats along

the main water channels, providing habitat for fish, amphibians, invertebrates, and a variety of bird species. These areas also act as natural filters, trapping sediments, heavy metals, and nutrients from runoff before they enter downstream rivers and lakes. The reed and sedge zones, often interspersed with shallow pools and seasonal water channels, support a variety of small mammals, waterbirds, and insect populations, which contribute to nutrient cycling and ecosystem resilience.

Riparian forests along the wetland edges are dominated by native tree species such as ***Ficus natalensis***, ***Markhamia lutea***, ***Phoenix reclinata***, and ***Syzygium guineense***.

These forested buffers provide critical ecosystem services, including bank stabilization, shade provision, and carbon sequestration. They serve as corridors for wildlife, connecting fragmented habitats and allowing species movement across the landscape. The forested patches are particularly important for bird species such as the Grey Crowned Crane (**Balearica regulorum**), African Jacana (**Actophilornis africanus**), herons, kingfishers, and a variety of migratory species that use the wetland as a stopover site.

Aquatic habitats within Nabakazi Wetland support diverse fish communities, which are critical for both ecological balance and local fisheries. Species such as tilapia (**Oreochromis niloticus**), catfish (**Clarias gariepinus**), and barbus species utilize vegetated areas for breeding, feeding, and shelter from predators. These fish populations contribute to nutrient

cycling by transferring energy between aquatic and terrestrial systems, and they provide an essential protein source for surrounding communities.

The wetland's ecological processes extend beyond species composition. Seasonal flooding replenishes soils with nutrients, maintains water levels during dry periods, and creates temporary habitats that support amphibians, insects, and migratory birds. Decomposition of organic material in the papyrus and reed beds contributes to carbon sequestration, making Nabakazi Wetland a valuable natural resource in the context of climate change mitigation. Its intact vegetation also mitigates local microclimatic extremes by cooling the air, maintaining humidity, and reducing the risk of drought stress for adjacent agricultural lands.



Despite its ecological richness, the wetland is under considerable threat from human activities. Encroachment into wetland zones for crop cultivation and settlement has reduced the extent of native vegetation. Invasive species, including water hyacinth (***Eichhornia crassipes***) and ***Lantana camara***, have displaced indigenous plants, disrupting habitat structure and ecological interactions. Additionally, unsustainable extraction of papyrus, wood, and other resources has degraded buffer zones, reducing the wetland's capacity to support biodiversity and ecosystem services. Loss of riparian forest cover has also diminished breeding and feeding habitats for birds and other wildlife, threatening the ecological integrity of the wetland.

The Nabakazi Wetland's ecological importance extends beyond local biodiversity. It serves as a critical component of Uganda's network of wetlands, contributing to regional water regulation, sediment control, and habitat connectivity. By sustaining fish breeding grounds, providing nesting sites for water birds, and maintaining hydrological balance,

the wetland plays a pivotal role in the ecological health of the Katonga, Kafu, Nkuse, and Muzizi river systems. Its conservation is therefore essential not only for protecting local species and livelihoods but also for maintaining ecosystem functions that underpin regional environmental stability and the sustainability of downstream freshwater systems, including Lake Victoria and the greater Nile Basin.

In conclusion, the Nabakazi Wetland represents a complex and highly valuable ecological system whose continued integrity depends on sustainable management and restoration. Its diverse habitats, dynamic ecological processes, and critical role in biodiversity conservation make it an indispensable natural asset for both people and wildlife. Effective restoration interventions must therefore integrate ecological recovery with sustainable livelihoods and community participation to ensure that the wetland continues to provide vital ecosystem services for current and future generations.

4.3 Socio-Economic Significance

The Nabakazi Wetland Ecosystem is not only ecologically important but also serves as a cornerstone for the livelihoods, food security, and socio-economic well-being of the communities living within and around its catchment area. Its resources and ecological services underpin a range of economic, cultural, and social activities that sustain tens of thousands of residents in Mubende, Kassanda, Gomba, Mpigi, and parts of Mityana districts.

Agriculture is the predominant livelihood activity in the wetland's surrounding subcounties. Communities rely on the nutrient-rich soils along the wetland margins for culti-

vating staple crops such as maize, beans, bananas, and sweet potatoes, as well as high-value cash crops like coffee and horticultural products. Seasonal flooding deposits fertile alluvial soils along riverbanks and floodplains, enhancing crop productivity without the need for synthetic fertilizers. Additionally, the wetland acts as a natural irrigation buffer, maintaining soil moisture and water availability during periods of low rainfall, thereby reducing the risk of crop failure and supporting household food security.

Fishing is another critical economic activity linked directly to the wetland's ecological integrity. The diverse fish populations sup-

ported by Nabakazi Wetland—such as tilapia, catfish, and barbus species—provide an essential source of protein for local households and contribute to the livelihoods of artisanal fishers. Seasonal breeding grounds within papyrus and reed zones are particularly important for sustaining fish stocks, which also support small-scale fish markets in nearby trading centers. The wetland thus plays a dual role in both nutrition and income generation for surrounding communities.

The wetland also supports a variety of non-timber resources that contribute to livelihoods and cultural practices. Papyrus is harvested for roofing, mat-making, and handicrafts, providing supplementary income to households, especially women and youth engaged in craft production. Riparian forests supply fuelwood, medicinal plants, and construction materials, which are essential for daily life in many households. Livestock farmers utilize wetland margins as grazing grounds during the dry season, taking advantage of the naturally regenerating grasses that provide feed for cattle, goats, and sheep.

Beyond direct economic benefits, the Nabakazi Wetland contributes to climate resilience and disaster risk reduction. Its capacity to regulate water flow mitigates flooding during rainy seasons and sustains water supply during dry periods, protecting agricultural lands, settlements, and infrastructure. Wetland vegetation reduces soil erosion and stabilizes riverbanks, safeguarding farmland and reducing sedimentation in downstream rivers and lakes. By maintaining microclimatic conditions, the wetland also reduces local temperature extremes and supports crop growth, which is particularly important under the increasing impacts of climate change.

The cultural and social significance of Nabakazi Wetland cannot be overlooked. Many communities hold traditional knowledge and practices related to wetland resource management, including taboos, customary fishing regulations, and sacred sites associated with rivers and forest patches. These cultural practices reinforce sustainable use patterns and contribute to biodiversity conservation.

However, the wetland's socio-economic benefits are increasingly threatened by unsustainable practices. Encroachment for agriculture and settlement has led to loss of buffer vegetation, while overharvesting of papyrus and firewood, coupled with weak enforcement of wetland regulations, has reduced the wetland's capacity to provide essential ecosystem services. Degradation of the wetland directly undermines livelihoods, reduces agricultural productivity, diminishes fish stocks, and increases vulnerability to floods and droughts. Communities dependent on the wetland therefore face both ecological and economic risks, highlighting the urgent need for structured restoration and sustainable management.

In conclusion, the Nabakazi Wetland is a vital socio-ecological system whose ecological health is directly linked to human well-being. Its role in sustaining livelihoods, enhancing food security, supporting cultural practices, and mitigating climate-related risks underscores the need for integrated conservation and restoration strategies. Protecting and rehabilitating the wetland will not only restore biodiversity and ecological processes but also safeguard the socio-economic stability and resilience of the communities that depend on it.

5.0 Justification for the Restoration Framework

The urgent need for a restoration framework for the Nabakazi Wetland Ecosystem arises from the combined pressures of ecological degradation, unsustainable resource use, and socio-economic dependence of surrounding communities on wetland resources. While the wetland provides vital ecosystem services that underpin regional water security, biodiversity, and livelihoods, its integrity has been increasingly compromised, threatening both environmental and human well-being.

5.1 Degradation Status and Drivers

Recent assessments, including the baseline and Grey crowned crested crane studies conducted by the National Association of Professional Environmentalists (NAPE), reveal that Nabakazi Wetland is under significant stress from human-induced and natural factors. Agricultural encroachment has been identified as the primary driver of degradation, with cultivation extending up to the highest water mark in many sections of the wetland. This has resulted in the clearing of natural vegetation, destabilization of riverbanks, and increased soil erosion. In addition, fuelwood collection, timber harvesting, and uncontrolled papyrus extraction have contributed to the loss of riparian forest cover and buffer vegetation, further reducing the wetland's resilience and capacity to provide ecological services.




The proliferation of invasive species, particularly water hyacinth (***Eichhornia crassipes***) and ***Lantana camara***, has disrupted native plant communities, altered hydrological dynamics, and reduced habitat quality for wildlife. Sedimentation resulting from upstream soil erosion has affected water quality and fish breeding

grounds, diminishing both biodiversity and the productivity of local fisheries. Human settlements along the wetland margins have increased waste discharge into wetland channels, exacerbating pollution and further threatening aquatic habitats.

Climate variability compounds these anthropogenic pressures. Erratic rainfall patterns, prolonged dry spells, and occasional floods intensify wetland degradation, reduce water availability for agriculture and domestic use, and threaten the reproductive cycles of key wetland species.

Collectively, these drivers undermine the wetland's capacity to sustain critical ecosystem services such as water regulation, soil fertility, carbon sequestration, and biodiversity support. Without deliberate restoration interventions, these trends are likely to escalate, resulting in long-term ecological, economic, and social consequences for the region.



5.2 Policy and Legal Justification

The restoration of Nabakazi Wetland is strongly supported by Uganda's national policies, legislation, and international obligations related to wetland conservation. At the national level, the Uganda Wetlands Policy (1995) emphasizes the sustainable utilization and rehabilitation of wetland resources to maintain their ecological functions and socio-economic benefits. The National Environment Act (2019) provides a legal framework for environmental management, including protection of wetlands, enforcement of buffer zones, and regulation of resource extraction activities. The Constitution of Uganda (1995, revised) further recognizes the state's responsibility to ensure sustainable management of natural resources for present and future generations.

On the regional and international front, Nabakazi Wetland falls within the scope of instruments that promote biodiversity conservation, water resource management, and

climate resilience. The Ramsar Convention (1971) identifies wetlands of international importance and encourages their wise use and restoration. The African-Eurasian Waterbird Agreement (AEWA) highlights the significance of wetlands as habitats for migratory waterbirds, while the Convention on Biological Diversity (CBD) underscores the importance of ecosystem restoration for maintaining species diversity and ecological processes. Additionally, the East African Community's policy frameworks on natural resources and climate adaptation reinforce the need for sustainable management of transboundary wetland systems that provide ecosystem services at regional scales.

From a practical standpoint, the restoration framework aligns with these policies and legal instruments by providing a structured, participatory, and scientifically informed approach to rehabilitate degrad-

ed areas, strengthen buffer zones, restore hydrological integrity, and enhance biodiversity. By integrating ecological restoration with community livelihoods, the framework not only fulfills regulatory requirements but also contributes to national and international targets on sustainable development, climate adaptation, and biodiversity conservation. It ensures that Nabakazi Wetland continues to provide critical ecosystem services, support resilient communities, and maintain its ecological role within the broader Nile Basin hydrological network.

The combined pressures of human activities, invasive species, and climate variability underscore the urgent need for a structured restoration framework. National and international policy instruments provide strong legal and institutional justification for interventions. By addressing the drivers of degradation and aligning restoration efforts with governance and livelihood objectives, the framework ensures the sustainable recovery of Nabakazi Wetland and the long-term provision of its ecological and socio-economic benefits.

6.0 Restoration Framework (Five Stages)

The restoration of Nabakazi Wetland requires a structured, multi-stage approach to ensure both ecological recovery and socio-economic benefits for the surrounding communities. Drawing on internationally recognized ecological restoration standards (SER, IUCN, FAO), the framework emphasizes adaptive management, stakeholder participation, and integration of ecological and livelihood interventions. The following five stages outline the practical steps for restoring Nabakazi Wetland.

6.1 Stage 1: Area Assessment

The initial stage focuses on understanding the current ecological, hydrological, and socio-economic conditions of the wetland. Comprehensive assessment involves a combination of field-based surveys, participatory consultations, and geospatial analysis.

Vegetation surveys document the extent of papyrus swamps, reed beds, riparian forests, and degraded areas. Aquatic surveys identify fish species composition, breeding habitats, and water quality parameters such as turbidity, pH, dissolved oxygen, and nutrient load. Soil assessments evaluate fertility, erosion patterns, and sediment deposition trends, while hydrological studies map water flow regimes, seasonal flood extents, and groundwater recharge areas.

Participatory methods, including focus group discussions, transect walks, and household interviews, provide insights into community resource use, dependency on wetland products, and perceptions of ecological degradation. These consultations are essential to identify both ecological priorities and social considerations for restoration planning.

Remote sensing and GIS mapping tools are used to produce spatially explicit maps of land use, wetland extent, degraded zones, and potential restoration sites. This baseline information forms the foundation for evidence-based decision-making, prioritization of interventions, and the design of monitoring indicators.

The outcome of this stage is a comprehensive assessment report that clearly identifies the causes and extent of degradation, existing ecological and socio-economic conditions, and opportunities for resto-

ration. This baseline ensures that subsequent restoration actions are targeted, feasible, and measurable.

6.2 Stage 2: Planning and Design

Building upon the assessment, the planning and design stage translates data into a structured restoration plan with defined objectives, strategies, and measurable outcomes.

Restoration goals are established based on ecological priorities, including re-vegetation of riparian buffers, restoration of fish breeding habitats, invasive species control, and enhancement of biodiversity. Socio-economic objectives are integrated, focusing on sustainable livelihoods, community participation, and climate resilience.

Specific intervention strategies are designed, including:

- **Natural regeneration:** Allowing degraded areas to recover through protective measures, such as exclusion of agricultural encroachment and livestock grazing.
- **Active reforestation and planting:** Planting indigenous trees, shrubs, and grasses in buffer zones to stabilize soils, provide wildlife habitat, and enhance carbon sequestration.
- **Hydrological restoration:** Constructing small check dams, contour trenches, or bunds to regulate water flow, reduce erosion, and improve groundwater recharge.
- **Invasive species management:** Systematic removal of water hyacinth and Lantana camara, coupled with reintroduction of native species to restore ecological balance.

Stakeholder consultation is central to this stage, ensuring that interventions are socially acceptable, technically feasible, and aligned with community priorities. Institu-

tional arrangements are defined, assigning clear roles and responsibilities to district authorities, NAPE technical teams, Community Wetland Management Committees, and partner organizations.

A detailed implementation schedule, budget estimates, and monitoring plan are developed to guide execution, with clearly defined success indicators for both ecological and socio-economic outcomes.

6.3 Stage 3: Implementation of Interventions

This stage involves executing the restoration strategies on the ground. Interventions are conducted in a coordinated manner, integrating ecological restoration with community development initiatives.

Ecological interventions include:

- Planting indigenous tree and shrub species along riverbanks and degraded buffer zones to prevent erosion and provide wildlife habitat.
- Establishing riparian corridors to enhance connectivity for birds, mammals, and aquatic species.
- Clearing invasive species and ensuring re-establishment of native vegetation to restore ecological function.
- Stabilizing soil through contour bunds, terraces, and sediment traps to reduce siltation of rivers and downstream lakes.

Socio-economic interventions run concurrently, enhancing community engagement and sustainable resource use.

These include:

- Promotion of alternative livelihoods such as beekeeping, fish farming, papyrus-based handicrafts, and small-scale horticulture.
- Training and capacity-building for community members on climate-smart agriculture, sustainable fishing practices, and wetland stewardship.

- Establishment of Community Wetland Management Committees (CWMCs) to oversee local monitoring, restoration maintenance, and compliance with wetland regulations.

Implementation is phased to allow adaptive learning, starting with priority degraded zones identified during the assessment stage and gradually scaling up to cover the entire wetland catchment. Regular stakeholder meetings ensure transparency, accountability, and alignment with ecological and socio-economic objectives.

6.4 Stage 4: Monitoring of Results

Monitoring is critical for assessing the effectiveness of restoration interventions, tracking progress, and guiding adaptive management. Monitoring combines scientific and community-based approaches to ensure comprehensive data collection.

Ecological indicators include:

- Area of wetland re-vegetated and extent of buffer zones restored.
- Water quality parameters, including turbidity, pH, nutrient levels, and dissolved oxygen.
- Species diversity, abundance, and presence of key indicator species, including the Grey Crowned Crane.
- Status of invasive species and success of eradication efforts.
- Socio-economic indicators include:
- Number of households benefiting from alternative livelihood initiatives.
- Adoption of sustainable farming and fishing practices.
- Community participation in wetland management and monitoring activities.

Monitoring is conducted quarterly and annually, with data used to produce progress reports and inform adaptive management decisions. GIS mapping and drone imagery are employed to track changes in vegetation cover and hydrological patterns, ensuring accurate and up-to-date information for decision-making.

drological patterns, ensuring accurate and up-to-date information for decision-making.

6.5 Stage 5:

Post-Restoration Management (Adaptive Management)

Long-term sustainability is achieved through adaptive management, which uses monitoring results to refine interventions and respond to changing ecological and socio-economic conditions.

The post-restoration stage includes:

- Continued management of restored vegetation, including periodic maintenance of planted areas and control of regrowth of invasive species.
- Strengthening institutional structures such as CWMCs to enforce wetland regulations and coordinate community engagement.
- Integration of wetland restoration into district development plans and local governance structures to ensure long-term support and funding.
- Exploration of financial sustainability mechanisms, including payments for ecosystem services, carbon credit initiatives, and eco-tourism opportunities.

Adaptive management ensures that restoration efforts remain relevant, resilient, and capable of maintaining the wetland's ecological integrity and socio-economic benefits over time. By embedding restoration into governance structures and local practices, the framework guarantees that Nabakazi Wetland continues to provide essential ecosystem services for generations to come.

The five-stage restoration framework provides a practical, scientifically informed, and participatory approach to

rehabilitate Nabakazi Wetland. By integrating ecological recovery with community livelihoods, adaptive management, and institutional support, the framework ensures that restoration efforts are effective, sustainable, and scalable.

7.0 Institutional Arrangements and Stakeholder Roles

The successful restoration and sustainable management of Nabakazi Wetland requires a coordinated, multi-level institutional framework that clearly defines the roles and responsibilities of all stakeholders involved. Given the wetland's ecological significance, socio-economic value, and trans-district reach, restoration efforts must integrate technical expertise, local governance structures, community participation, and multi-sectoral collaboration. The institutional arrangements outlined below are designed to ensure efficiency, accountability, and sustainability.

7.1 District and Local Government Authorities

District authorities, particularly the Environmental and Natural Resources Offices, play a central role in facilitating wetland restoration activities.

Their responsibilities include:

- Providing technical guidance and oversight during the planning, implementation, and monitoring phases of restoration.
- Ensuring alignment of restoration interventions with district development plans, environmental policies, and legal frameworks, including the National Environment Act (2019) and Uganda's Wetlands Policy (1995).
- Coordinating resource mobilization, in-

cluding allocation of budgetary support for community engagement, reforestation activities, and monitoring programs.

- Enforcing wetland protection regulations, preventing encroachment, and addressing non-compliance by community members or commercial actors.

Sub-county authorities serve as the primary link between district offices and local communities. They facilitate participatory planning, mobilize community members for restoration activities, and monitor compliance with agreed restoration protocols at the village level.

7.2 National Association of Professional Environmentalists (NAPE) Technical Team

The NAPE technical team provides specialized expertise in ecological restoration, biodiversity management, capacity building, media advocacy, outreach and environmental monitoring.

Their responsibilities include:

- Conducting ecological and hydrological assessments to inform evidence-based restoration planning.
- Designing restoration interventions, including native species planting, invasive species management, and hydrological rehabilitation.
- Training local stakeholders, including district staff, community members, and school groups, on best practices for wetland restoration and sustainable resource use.
- Communication and publicity.
- Monitoring and evaluating ecological recovery and providing adaptive management recommendations based on data collected.

NAPE and Likeminded groups role is critical in ensuring that restoration activities are scientifically sound, technically feasible, and

aligned with local ,national ,regional and international ecological restoration standards.

7.3 Community Wetland Management Committees (CWMCs)

CWMCs are community-led bodies established to manage local wetland resources, coordinate restoration activities, and promote sustainable use.

Their responsibilities include:

- Mobilizing local participation in planting, invasive species control, and other on-the-ground restoration efforts.
- Monitoring wetland conditions, reporting illegal encroachment or unsustainable resource extraction, and liaising with district authorities.
- Facilitating knowledge sharing and training sessions on sustainable agriculture, fishing practices, and alternative livelihoods.
- Supporting local enforcement of agreed wetland buffer zones and conservation rules, thereby fostering a sense of ownership and stewardship among community members.

By placing communities at the center of wetland management, CWMCs enhance long-term sustainability and reduce conflicts over resource use.

7.4 Government Agencies and Sectoral Partners

Several government institutions and sectoral agencies provide complementary support for wetland restoration.

Key roles include:

- **Ministry of Water and Environment (MWE):** Policy oversight, technical guidance, and monitoring compliance with wetland protection regulations.

- **National Forestry Authority (NFA):** Provision of tree seedlings, technical advice on reforestation, and enforcement against illegal logging in wetland buffer zones.
- **Fisheries Department:** Guidance on sustainable fishing practices, fish breeding habitat restoration, and monitoring aquatic biodiversity.
- **Local Development Agencies:** Support in integrating restoration activities into broader socio-economic development initiatives, such as climate-smart agriculture and livelihood diversification programs.

7.5 Private Sector and Civil Society Partners

The private sector and civil society organizations play an important role in supporting restoration through funding, technical inputs, and community engagement.

Responsibilities include:

- Co-financing restoration activities, such as tree planting, alternative livelihood projects, and water infrastructure development.
- Providing technical expertise, innovation, and best practices from other successful wetland or watershed restoration initiatives.
- Engaging in awareness campaigns, community education, and capacity-building initiatives to promote sustainable resource use.
- Collaborating with district authorities and communities to implement climate-smart practices and explore eco-tourism or carbon credit opportunities.
- Providing both finance and kind support to support restoration practice.

7.6 Coordination Mechanisms

Effective coordination among all stakeholders is essential for the success of the Nabakazi Wetland Restoration Framework.

This is achieved through:

- **Technical Steering Committees:** Comprising district officials, NAPE experts, sectoral agencies, and CWMC representatives to oversee planning, implementation, and evaluation.

- **Regular Stakeholder Meetings:** Quarterly or biannual meetings to review progress, address challenges, and share lessons learned.
- **Reporting and Accountability Frameworks:** Clear reporting lines between CWMCs, district authorities, and technical teams to ensure transparency and timely decision-making.

By fostering collaboration and clearly defining roles, the institutional framework ensures that restoration interventions are coherent, inclusive, and sustainable.

The institutional arrangements for Nabakazi Wetland Restoration provide a robust governance structure that integrates technical expertise, local stewardship, and multi-sectoral collaboration. By assigning clear roles and responsibilities, fostering community ownership, and promoting effective coordination, these arrangements create an enabling environment for sustainable wetland management and long-term ecological recovery.

8.0 Implementation Strategy and Operational Plan

The successful restoration of Nabakazi Wetland depends on a well-structured, phased, and practical implementation strategy that integrates ecological, socio-economic, and governance considerations. This operational plan provides a roadmap for executing the five-stage restoration framework, ensuring that interventions are effective, participatory, and sustainable.

8.1 Phased Implementation Approach

Restoration interventions are implemented in a phased manner to allow for adaptive learning, effective resource allocation, and community engagement. The implementation is organized into three main phases:

Phase 1: Priority Area Restoration

The first phase focuses on highly degraded zones and ecologically sensitive areas, such as riverbanks, fish breeding grounds,

and riparian forest fragments.

Activities include:

- Removal of invasive plant species, including water hyacinth and *Lantana camara*.
- Establishment of buffer zones by stone marking them, planting indigenous trees, shrubs, and grasses.
- Soil stabilization measures, including contour bunds, terraces, and sediment traps.
- Community sensitization and mobilization for participation in restoration activities.

Phase 2: Scaling Up to Wider Wetland Zones

Once priority areas have been stabilized and restoration practices tested, interventions are expanded to cover broader sections of the wetland catchment beyond the 4 sub counties of mubende.

Activities include:

- Reforestation of degraded upland areas to enhance watershed protection.
- Hydrological restoration through small water retention structures, improved drainage channels, and floodplain management.
- Protection and recovery of degraded breeding sites for bird species native to the wetland.
- Promotion of alternative livelihoods, including fish farming, beekeeping, and papyrus handicraft production, to reduce pressure on wetland resources.

Phase 3: Consolidation and Sustainability

The final phase ensures that restoration gains are maintained and scaled for long-term impact.

Activities include:

- Strengthening community-based wetland management committees (CWMCs) for ongoing monitoring and enforcement.
- Capacity-building and training programs for local farmers, fishers, and youth in sustainable resource use and climate-smart practices.
- Integration of restored wetlands into local development plans and climate adaptation strategies.
- Establishment of financial sustainability

ty mechanisms, including payments for ecosystem services, carbon credits, and eco-tourism initiatives.

8.2 Community Engagement and Participation

Community involvement is central to the operational plan, recognizing that sustainable restoration cannot be achieved without active local participation. Engagement strategies include:

- **Participatory Planning Workshops:** Communities are consulted to identify local priorities, select restoration sites, and define acceptable resource-use rules.
- **Awareness Campaigns:** Education on wetland ecosystem services, climate change impacts, and sustainable livelihoods enhances community buy-in.
- **Volunteer Programs:** Local residents, schools, and youth groups participate in tree planting, invasive species removal, and monitoring activities.
- **Livelihood Support:** Introduction of alternative income-generating activities reduces dependency on wetland resources and strengthens community resilience.

Through these mechanisms, communities develop a sense of ownership, ensuring that restored areas are protected and maintained over the long term.

8.3 Resource Mobilization and Management

Effective implementation requires careful planning and mobilization of financial, human, and material resources. Key elements include:

- **Financial Resources:** Funding from district budgets, national government programs, development partners, NGOs, donors and private sector contributions.
- **Technical Expertise:** NAPE technical team, district local government, and sectoral agencies provide guidance on ecological restoration, hydrology, and biodiversity management.
- **Material Resources:** Seedlings, planting tools, soil stabilization materials, and monitoring equipment are procured and distributed

ed according to the implementation plan.

- **Human Resources:** Community mobilizers, volunteers, and trained CWMC members provide labor and support for on-the-ground activities.

Transparent financial and resource management systems are established to ensure accountability, efficiency, and timely disbursement of funds and materials.

8.4 Timelines and Work Plan

The operational plan is designed for a five-year restoration cycle, with specific milestones aligned to ecological and socio-economic outcomes:

- **Year 1:** Baseline assessments, participatory planning, initial community mobilization, and restoration of priority degraded zones.
- **Year 2–3:** Expansion of restoration interventions, reforestation of buffer zones, invasive species management, and establishment of alternative livelihoods.
- **Year 4:** Consolidation of restored areas, scaling up hydrological interventions, and community capacity-building.
- **Year 5:** Final adaptive management, institutionalization of CWMCs, integration into district development plans, and evaluation of ecological and socio-economic outcomes.

Timelines are flexible to accommodate adaptive management, climatic variability, and community engagement dynamics, ensuring that restoration remains responsive to emerging challenges.

8.5 Monitoring, Evaluation, and Learning

Monitoring and evaluation (MEL) are embedded within the operational plan to track progress, measure outcomes, and inform adaptive management:

- **Ecological Monitoring:** Vegetation recovery, water quality, species diversity, and hydrological performance are tracked regularly using scientific and community-based methods.

- **Socio-Economic Monitoring:** Household livelihoods, adoption of sustainable practices, and community participation levels are evaluated to assess social impacts.
- **Learning and Feedback:** Data from monitoring activities inform adjustments to restoration techniques, stakeholder engagement strategies, and resource allocation.

Regular reporting, feedback loops, and stakeholder review meetings ensure transparency, accountability, and continuous improvement.

8.6. IMPLEMENTATION SCHEDULE AND RESOURCE PLAN

8.6.1 Implementation Approach

The implementation strategy is based on four complementary pillars:

- **Ecological Restoration:** Physical rehabilitation of degraded wetland and buffer areas.
- **Livelihood Enhancement:** Support for sustainable, nature-based income-generating activities.
- **Institutional Strengthening:** Building local governance capacity for effective wetland management.
- **Monitoring and Adaptive Learning:** Continuous assessment of progress and adjustment of interventions as needed.

Each pillar is implemented through an integrated, multi-stakeholder process involving the Mubende District Environment Office, NAPE technical team, Community Wetland Management Committees (CWMCs), and partner organizations.

8.6.2 Phased Implementation Plan (2025–2030)

The framework will be implemented in three main phases:

Phase I: Preparatory and Foundational Activities (Year 1)

- Establishment of coordination structures (NW-TSC, CWMCs, TWGs).
- Detailed site-level assessments and mapping of degraded areas.

- Boundary demarcation and buffer zone identification.
- Community sensitization and training on wetland laws and restoration techniques.
- Initiation of pilot restoration sites in Madudu and Butologo subcounties.
- Establishment of indigenous plant nurseries and papyrus regeneration plots.

Expected Outputs:

- Functional institutional structures and governance frameworks.
- Detailed wetland maps and baseline monitoring database.
- 100 hectares of priority restoration initiated.
- At least 300 community members trained in restoration and monitoring.

Phase II:

Full-Scale Implementation (Years 2–3)

- Expansion of restoration activities to degraded areas in Kitenga, Kalonga, and Bwakago.
- Active reforestation of riparian zones and hydrological rehabilitation (check dams, sediment traps).
- Removal of invasive species (Lantana camara, water hyacinth) and reintroduction of native vegetation.
- Promotion of sustainable livelihood enterprises (apiary, aquaculture, craft production).
- Integration of restoration activities into sub-county development and land-use plans.
- Strengthening of CWMCs for local monitoring and enforcement.

Expected Outputs:

- 500 hectares of restored wetland and riparian habitat.
- At least 400 households engaged in alternative livelihoods.
- Reduced wetland encroachment and siltation rates.
- Functional local enforcement and monitoring committees in all subcounties.

Phase III: Consolidation, Monitoring, and Sustainability (Years 4–5)

- Maintenance of restored sites and replanting in areas with low survival rates.
- Continued support for community livelihood groups and cooperatives.
- Integration of restoration lessons into district and national policy frameworks.
- Establishment of long-term financing mechanisms (carbon credits, eco-tourism, PES).
- Comprehensive ecological and socio-economic evaluation to assess progress toward targets.

Expected Outputs:

- Over 1,000 hectares of wetland and buffer zones restored and protected.
- 60% increase in vegetation cover and return of indicator species.
- CWMCs integrated into sub-county governance structures.
- Evidence-based adaptive management plan for post-2030 continuation.

8.6. 3 Indicative Five-Year Work Plan (Narrative Summary)

Year	Focus	Major Outputs	Key Actors
2025	Preparatory phase: baseline surveys, boundary demarcation, and institutional setup.	Governance structures established, pilot sites launched, nurseries set up.	NAPE, DEO, CWMCs, Local Councils
2026	Expansion phase I: large-scale planting, hydrological restoration, community mobilization.	300+ ha restored, check dams constructed, 10 community groups trained.	DEO, NAPE, MWE, Communities
2027	Expansion phase II: invasive species control, livelihood integration, awareness campaigns.	500+ households trained, 600 ha restored, increased enforcement compliance.	CWMCs, NGOs, Local Leaders
2028	Consolidation: replanting, monitoring, ecosystem stabilization.	Ecological stability achieved in 80% of restoration zones.	NAPE, DEO, CWMCs
2029–2030	Sustainability and scaling: adaptive management, financing, policy mainstreaming.	Sustainable financing initiated, 1,000+ ha restored, policy integration complete.	MWE, NAPE, Partners

8.6.4 Resource Requirements and Indicative Budget Categories

The funding strategy combines multiple streams to ensure long-term financial sustainability:

- **Public Funding:** District and sub-county environment budgets, supplemented by the Ministry of Water and Environment.
- **Development Partner Contributions:** Ongoing collaboration with NAPE and international partners such as Partner für Klima und Natur (Germany) and WWF Uganda.
- **Private Sector and CSR Initiatives:** Partnerships with coffee cooperatives, agro-businesses, and local enterprises investing in green production.
- **Community Contributions:** In-kind labor, nursery management, and maintenance of restoration sites.
- **Innovative Financing Mechanisms:** Development of payment for ecosystem services (PES), carbon credit schemes, and eco-tourism ventures.

While specific cost figures will be refined during project budgeting, the indicative resource requirements are organized under key thematic categories:

Budget Category	Indicative Activities	Main Implementers
1. Technical Coordination and Capacity Building	Establishment of NW-TSC and CWMCs, training workshops, staff facilitation.	DEO, NAPE, MWE
2. Ecological Restoration	Seedling production, tree planting, hydrological rehabilitation structures, invasive species removal.	DEO, NAPE, CWMCs, Communities and other CSO groups.
3. Livelihood and Community Empowerment	Apiary kits, aquaculture setup, craft training, micro-credit support.	NGOs, CWMCs, District CDO
4. Monitoring and Evaluation	Ecological surveys, GIS mapping, water quality monitoring, socio-economic assessments.	NAPE, DEO, CWMCs
5. Awareness and Education	Radio programs, school campaigns, IEC materials, Wetlands Day events.	DEO, Local Councils, NAPE, other partner CSOs Media
6. Logistics and Equipment	Field transport, tools, GPS and water testing kits, protective gear.	District Administration
7. Sustainability and Financing Mechanisms	Feasibility studies for PES/carbon credits, eco-tourism development.	NAPE, Partners, Private Sector

Indicative Total Cost:

While precise figures will depend on donor commitments and district allocations, a preliminary estimate suggests **a total investment requirement of USD 1.2–1.5 million** over five years to achieve full-scale restoration and sustainability outcomes

8.6 Implementation Oversight and Accountability

A robust accountability mechanism ensures transparency and efficiency throughout implementation:

- **Quarterly Technical Review Meetings** at district level to track progress and expenditures.
- **Annual Audits and Performance Reviews** conducted jointly by NAPE, DEO, and external auditors.
- **Community Scorecards** used to assess satisfaction, participation, and benefits at local levels.

- **Public Reporting:** Annual progress summaries disseminated through community radio and district websites.

The five-year implementation and resource plan provides a structured and achievable pathway for the restoration of Nabakazi Wetland. Through phased action, strategic partnerships, and transparent resource management, the plan ensures that both ecological recovery and community benefits are realized sustainably.

8.7 Risk Management

The operational plan identifies and mitigates potential risks, including:

- **Environmental Risks:** Floods, droughts, and invasive species outbreaks are managed through adaptive restoration practices and early warning systems.

- **Socio-Economic Risks:** Community conflicts, overexploitation of resources, and lack of participation are addressed through inclusive engagement, awareness campaigns, and livelihood support.
- **Institutional Risks:** Weak coordination and inadequate technical capacity are mitigated through defined roles, capacity-building, and multi-stakeholder governance mechanisms.

The implementation strategy and operational plan provide a comprehensive roadmap for restoring Nabakazi Wetland in a structured, participatory, and adaptive manner. By integrating ecological interventions with community engagement, resource management, and institutional support, the plan ensures that restoration efforts are sustainable, measurable, and capable of delivering long-term ecological and socio-economic benefits.

9. Monitoring, Evaluation, and Adaptive Management

Monitoring, Evaluation, and Adaptive Management (MEAM) are critical components of the Nabakazi Wetland Restoration Framework, ensuring that restoration interventions achieve their ecological and socio-economic objectives while remaining responsive to changing conditions. By systematically tracking progress, evaluating outcomes, and applying lessons learned, MEAM strengthens decision-making, accountability, and sustainability of wetland restoration efforts.

9.1 Objectives of Monitoring and Evaluation

The primary objectives of MEAM in the Nabakazi Wetland context are to:

- **Assess Ecological Recovery:** Monitor the restoration of native vegetation, riparian buffer zones, and aquatic habitats to ensure that ecosystem functions such as sediment filtration, water regulation, and biodiversity support are restored.
- **Measure Socio-Economic Impacts:** Evaluate the extent to which restoration interventions improve community livelihoods, food security, and resilience to climate variability.
- **Track Implementation Progress:** Ensure that restoration activities, as outlined in the operational plan, are executed on schedule, within budget, and according to technical standards.
- **Inform Adaptive Management:** Provide data-driven insights to modify restoration techniques, community engagement approaches, and governance mechanisms in response to observed outcomes and emerging challenges.

9.2 Monitoring Indicators

Monitoring combines ecological, hydrological, and socio-economic indicators, structured around measurable, achievable, and time-bound targets.

Ecological Indicators:

- **Vegetation Recovery:** Extent of reforested buffer zones, survival rate of planted indigenous trees and grasses, and natural regeneration of wetland plants.
- **Biodiversity:** Species richness and abundance of key indicator species, including birds (e.g., Grey Crowned Crane), fish, amphibians, and pollinators.
- **Invasive Species Control:** Reduction in coverage of invasive species such as water hyacinth and Lantana camara.
- **Hydrological Function:** Changes in water levels, flow rates, and sedimentation patterns within the wetland and downstream catchments.

9.2 Monitoring Indicators

Monitoring combines ecological, hydrological, and socio-economic indicators, structured around measurable, achievable, and time-bound targets.

Ecological Indicators:

- **Vegetation Recovery:** Extent of reforested buffer zones, survival rate of planted indigenous trees and grasses, and natural regeneration of wetland plants.
- **Biodiversity:** Species richness and abundance of key indicator species, including birds (e.g., Grey Crowned Crane), fish, amphibians, and pollinators.

dance of key indicator species, including birds (e.g., Grey Crowned Crane), fish, amphibians, and pollinators.

- **Invasive Species Control:** Reduction in coverage of invasive species such as water hyacinth and Lantana camara.
- **Hydrological Function:** Changes in water levels, flow rates, and sedimentation patterns within the wetland and downstream catchments.

Indicator	Method / Tool	Frequency	Responsible Entity
Vegetation cover recovery (ha)	GIS and field transects	Bi-annual	DLG, NAPE
Species diversity (flora and fauna)	Biodiversity surveys	Annual	NAPE, Nature Uganda
Water quality (turbidity, pH, nutrients)	Water testing kits	Quarterly	DLG, MWE, CWMCs
Sediment load and siltation rate	Riverbank sampling	Annual	NAPE, MWE
Groundwater levels and flow regime	Observation wells, hydrological mapping	Quarterly	DLG, NEMA, MWE

Socio-Economic Indicators:

- **Household Livelihoods:** Number of households adopting alternative income-generating activities, including beekeeping, fish farming, and sustainable papyrus use.
- **Food Security:** Availability and diversity of agricultural and fishery products resulting from wetland restoration.
- **Community Participation:** Engagement in restoration activities, membership in Community Wetland Management Committees (CWMCs), and compliance with wetland protection rules.
- **Awareness and Knowledge:** Increase in community understanding of wetland functions, climate resilience, and sustainable resource management practices.

Indicator	Method / Tool	Frequency	Responsible Entity
Number of households with alternative livelihoods	Household surveys	Annual	CWMCs, CDO
Income increase from sustainable enterprises (%)	Socio-economic surveys	Annual	DEO, NAPE
Reduction in illegal encroachment incidents	Compliance records	Quarterly	CWMCs, Local Councils
Food security levels among wetland households	Nutrition and income surveys	Bi-annual	District Planning Unit
Participation of women and youth in restoration	Attendance and membership records	Continuous	CWMCs, NAPE

9.3 Monitoring Methods

A combination of scientific and community-based monitoring methods ensures comprehensive data collection:

- **Field Surveys:** Vegetation plots, bird point counts, fish sampling, and soil assessments provide quantitative ecological data.
- **Remote Sensing and GIS Mapping:** Satellite imagery and drone surveys track changes in wetland extent, vegetation cover, and habitat connectivity over time.
- **Community-Based Monitoring:** CWMCs, local volunteers, and school groups participate in observation and data recording, enhancing ownership and local knowledge.
- **Water Quality Testing:** Regular measurement of parameters such as turbidity, dissolved oxygen, pH, and nutrient load ensures that hydrological improvements are sustained.
- **Household Surveys and Focus Group Discussions:** Assess changes in livelihoods, food security, and community perceptions of restoration effectiveness.

9.4 Evaluation and Reporting

Evaluation involves analyzing monitoring data to determine whether restoration objectives are being met. This includes:

- Comparing ecological and socio-economic indicators against baseline data established in Stage 1.
- Identifying areas of success, challenges, and gaps in implementation.
- Producing quarterly and annual progress reports that document restoration outcomes, lessons learned, and recommendations for improvement.
- Sharing evaluation results with all stakeholders, including district authorities, NAPE

technical teams, CWMCs, and partner organizations, to ensure transparency and informed decision-making.


9.5 Adaptive Management

Adaptive management is an iterative process where monitoring and evaluation results are used to refine restoration interventions.

Key elements include:

- **Feedback Loops:** Data collected from ecological and socio-economic monitoring informs adjustments to planting strategies, invasive species control, and community engagement approaches.
- **Flexible Planning:** Restoration techniques are modified based on observed outcomes, environmental conditions, and community feedback. For example, planting schedules may be adjusted to align with rainfall patterns, or new species may be introduced to enhance biodiversity.
- **Capacity Building:** Continuous training of district staff, technical teams, and community members ensures that adaptive practices are effectively implemented and knowledge is retained locally.
- **Policy Integration:** Insights from monitoring inform district and national policy development, reinforcing legal compliance and promoting broader wetland conservation objectives.

MONITORING, EVALUATION, AND ADAPTIVE MANAGEMENT (MEAM))



Through adaptive management, the Nabakazi Wetland Restoration Framework remains resilient to environmental variability, social dynamics, and unforeseen challenges, ensuring that both ecological recovery and community benefits are sustained over the long term.

Monitoring, Evaluation, and Adaptive Management provide a structured, evidence-based approach to track progress, measure impacts, and enhance the effectiveness of restoration interventions. By combining scientific rigor with community participation, MEAM ensures that the Nabakazi Wetland continues to deliver essential ecosystem services, support resilient livelihoods, and maintain biodiversity. The iterative nature of adaptive management guarantees that restoration efforts remain responsive, sustainable, and capable of achieving long-term ecological and socio-economic objectives.

10. Sustainability Measures and Long-Term Management

Ensuring the long-term success of the Nabakazi Wetland restoration requires a holistic approach that integrates ecological, socio-economic, institutional, and financial sustainability. Without

deliberate sustainability measures, gains achieved through restoration may be lost due to renewed human pressures, environmental variability, or institutional weaknesses. This section outlines key strategies to maintain and enhance the wetland's ecological integrity while supporting community livelihoods over the long term.

10.1 Ecological Sustainability

Ecological sustainability focuses on maintaining the health, functionality, and resilience of the wetland ecosystem.

Key strategies include:

- **Protection of Restored Areas:** Implementing and enforcing buffer zones along riverbanks, riparian forests, and wetland margins prevents encroachment and preserves habitats. Signage, demarcation, and community patrols support long-term protection.
- **Biodiversity Conservation:** Continuous monitoring of flora and fauna ensures the survival of key species, particularly indicator and threatened

species such as the Grey Crowned Crane. Restoration of native vegetation, wetlands, and aquatic habitats enhances ecological connectivity and resilience.

- **Invasive Species Management:** Sustained removal of invasive plants, coupled with reintroduction of native species, prevents ecological imbalance and supports natural regeneration.
- **Hydrological Integrity:** Maintaining natural water flow regimes, protecting water channels, and controlling sedimentation ensures that the wetland continues to provide essential services such as flood regulation, groundwater recharge, and water purification.

By maintaining these ecological interventions, the wetland can continue to provide ecosystem services critical for regional biodiversity, agriculture, and climate resilience.

10.2 Community and Livelihood Sustainability

The social dimension of sustainability emphasizes the continued participation of local communities and the development of resilient livelihoods.

Key strategies include:

- **Community Ownership and Stewardship:** CWMCs and other community governance structures are strengthened to oversee restoration, enforce wetland regulations, and coordinate local monitoring activities. Community participation fosters a sense of responsibility and ensures adherence to sustainable resource use practices.
- **Alternative Livelihoods:** Promoting income-generating activities that do not degrade the wetland, such as bee-keeping, sustainable papyrus craft production, fish farming, and eco-tourism, reduces dependency on extractive activities while enhancing household income.
- **Capacity Building:** Ongoing training and knowledge-sharing programs build community skills in climate-smart agri-

culture, wetland management, and disaster risk reduction. This ensures that residents are equipped to manage resources sustainably and adapt to environmental changes.

- **Education and Awareness:** Community sensitization campaigns, school programs, and public outreach strengthen understanding of wetland ecosystem services and the importance of conservation, ensuring cultural and behavioral reinforcement of sustainable practices.

Through these measures, communities become active partners in wetland conservation, guaranteeing the long-term social and economic sustainability of restoration interventions.

10.3 Institutional and Governance Sustainability

Institutional sustainability ensures that restoration interventions are embedded within functional governance structures.

Strategies include:

- **Integration into District Plans:** Incorporating wetland restoration and management into district and sub-county development plans ensures political and administrative support, as well as long-term funding allocation.
- **Multi-Stakeholder Coordination:** Strengthening coordination among district authorities, NAPE technical teams, CWMCs, government agencies, private sector actors, and NGOs ensures continuity, avoids duplication, and promotes shared responsibility.
- **Legal Compliance and Enforcement:** Enforcement of existing wetland policies, regulations, and buffer zone requirements supports long-term protection, while legal frameworks provide avenues for addressing violations.
- **Institutional Capacity Building:** Continuous training for district officers, CWMCs, and technical staff ensures that skills and knowledge required for restoration management are retained locally and can be transferred to future practitioners.

- Institutional sustainability safeguards the framework from governance gaps and ensures that restoration remains a priority within local and national environmental agendas.

10.4 Financial Sustainability

Long-term restoration requires stable and diversified funding sources.

Financial sustainability measures include:

- **Government and Donor Funding:** Continued allocation of funds from district budgets, national government programs, and development partners to support core restoration activities.
- **Private Sector Partnerships:** Engagement with coffee cooperatives, agro-industrial companies, and other private actors provides additional investment in wetland restoration and sustainable livelihoods.
- **Payments for Ecosystem Services (PES):** Initiatives such as carbon credits, water quality improvement payments, and biodiversity offsets provide incentives for maintaining wetland integrity while generating revenue.
- **Eco-Tourism Development:** Sustainable tourism opportunities, such as bird-watching for the Grey Crowned Crane and guided wetland tours, generate income for communities and local management bodies.
- **Community Contribution Mechanisms:** Participatory approaches, including voluntary labor, local fundraising, and cooperative management of restoration enterprises, supplement financial resources and strengthen ownership.

Diversifying financial sources ensures that restoration activities can continue beyond the initial project cycle, sustaining ecological and socio-economic benefits for future generations.

10.5 Knowledge Management and Learning

Sustainability is reinforced through systematic documentation and learning:

- **Restoration Records:** Maintaining detailed records of restoration activities, monitoring data, and outcomes provides institutional memory and facilitates replication.
- **Research and Innovation:** Encouraging applied research on wetland management, climate adaptation, and sustainable livelihoods enhances evidence-based interventions.
- **Knowledge Sharing Platforms:** Workshops, community forums, and digital repositories disseminate lessons learned, best practices, and innovations, fostering continuous improvement and adaptation.

By embedding knowledge management into long-term strategies, Nabakazi Wetland restoration remains adaptive, resilient, and scalable.

Long-term sustainability of Nabakazi Wetland restoration depends on an integrated approach that combines ecological integrity, community engagement, institutional support, financial resilience, and knowledge management. By aligning restoration interventions with social, economic, and governance frameworks, the wetland can continue to provide essential ecosystem services, support resilient livelihoods, and contribute to regional biodiversity and climate adaptation objectives. These measures ensure that restoration gains are maintained, scaled, and adapted to evolving environmental and socio-economic challenges, securing the wetland's value for present and future generations.

11. References

- National Association of Professional Environmentalists (NAPE). (2025). *Baseline Study Report on Potential Sites for Restoration along Nabakazi Wetland Ecosystem in Mubende District.*
- National Association of Professional Environmentalists (NAPE) (2025/6) *Ecological Study and Mapping of Breeding sites of the Grey Crowned Crane in Mubende, Kasanda and Gomba Districts-Uganda*
- National Association of Professional Environmentalists (NAPE) (2025/6) –*Quarterly restoration pilot field Progress reports*
- Society for Ecological Restoration (SER), IUCN, FAO. (2019). *International Standards for the Practice of Ecological Restoration.*
- National Environment Management Act (2019), Uganda.
- Uganda Wetlands Policy (1995).
- Ramsar Convention Secretariat. (2018). *Ramsar Handbook on Wise Use of Wetlands.*
- African-Eurasian Waterbird Agreement (AEWA).
- Constitution of the Republic of Uganda (1995, revised 2017).





A publication of:
**National Association of Professional Environmentalists
(NAPE)**

With support from:
Partner für Klima und Natur gGmbH (PKN).



**PARTNER FÜR
KLIMA UND NATUR**